

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1626GMS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 DEC 21 IPC search and display fields enhanced in CA/CAPLUS with the
IPC reform
NEWS 4 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/
USPAT2
NEWS 5 JAN 13 IPC 8 searching in IFIPAT, IFIUDB, and IFICDB
NEWS 6 JAN 13 New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to
INPADOC
NEWS 7 JAN 17 Pre-1988 INPI data added to MARPAT
NEWS 8 JAN 17 IPC 8 in the WPI family of databases including WPIFV
NEWS 9 JAN 30 Saved answer limit increased
NEWS 10 JAN 31 Monthly current-awareness alert (SDI) frequency
added to TULSA
NEWS 11 FEB 21 STN AnaVist, Version 1.1, lets you share your STN AnaVist
visualization results
NEWS 12 FEB 22 Status of current WO (PCT) information on STN
NEWS 13 FEB 22 The IPC thesaurus added to additional patent databases on STN
NEWS 14 FEB 22 Updates in EPFULL; IPC 8 enhancements added
NEWS 15 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 16 FEB 28 MEDLINE/LMEDLINE reload improves functionality
NEWS 17 FEB 28 TOXCENTER reloaded with enhancements
NEWS 18 FEB 28 REGISTRY/ZREGISTRY enhanced with more experimental spectral
property data
NEWS 19 MAR 01 INSPEC reloaded and enhanced
NEWS 20 MAR 03 Updates in PATDPA; addition of IPC 8 data without attributes
NEWS 21 MAR 08 X.25 communication option no longer available after June 2006
NEWS 22 MAR 22 EMBASE is now updated on a daily basis
NEWS 23 APR 03 New IPC 8 fields and IPC thesaurus added to PATDPAFULL
NEWS 24 APR 03 Bibliographic data updates resume; new IPC 8 fields and IPC
thesaurus added in PCTFULL
NEWS 25 APR 04 STN AnaVist \$500 visualization usage credit offered

NEWS EXPRESS FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.
V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT
<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that

04/06/2006 10717786.trn

specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:37:04 ON 06 APR 2006

=>

Uploading

THIS COMMAND NOT AVAILABLE IN THE CURRENT FILE

Do you want to switch to the Registry File?

Choice (Y/n):

Switching to the Registry File...

Some commands only work in certain files. For example, the EXPAND command can only be used to look at the index in a file which has an index. Enter "HELP COMMANDS" at an arrow prompt (=>) for a list of commands which can be used in this file.

=> FILE REGISTRY

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 09:37:13 ON 06 APR 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 4 APR 2006 HIGHEST RN 879269-14-4

DICTIONARY FILE UPDATES: 4 APR 2006 HIGHEST RN 879269-14-4

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.


04/06/2006 10717786.trn

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10717786.str



chain nodes :

6 7 8

ring nodes :

1 2 3 4 5 9 10 11 12 13 14 15 16 17 18 19 20

chain bonds :

1-6 2-7 5-20 7-8 8-10

ring bonds :

1-2 1-5 2-3 3-4 4-5 9-10 9-14 10-11 11-12 12-13 13-14 15-16 15-20
16-17 17-18 18-19 19-20

exact/norm bonds :

1-2 1-5 1-6 2-3 2-7 4-5 5-20 7-8 8-10

exact bonds :

3-4

normalized bonds :

9-10 9-14 10-11 11-12 12-13 13-14 15-16 15-20 16-17 17-18 18-19 19-20

isolated ring systems :

containing 1 : 9 : 15 :

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:CLASS 7:CLASS 8:CLASS 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom
20:Atom

L1 STRUCTURE UPLOADED

=> d 11

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 09:37:28 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 158 TO ITERATE

100.0% PROCESSED 158 ITERATIONS 7 ANSWERS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 2406 TO 3914
PROJECTED ANSWERS: 7 TO 298

L2 7 SEA SSS SAM L1

=> s l1 sss full

FULL SEARCH INITIATED 09:37:35 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 3528 TO ITERATE

100.0% PROCESSED 3528 ITERATIONS
SEARCH TIME: 00.00.01

132 ANSWERS

L3 132 SEA SSS FUL L1

=> FIL HCAPLUS

| COST IN U.S. DOLLARS | SINCE FILE ENTRY | TOTAL SESSION |
|----------------------|------------------|---------------|
| FULL ESTIMATED COST | 167.38 | 167.59 |

FILE 'HCAPLUS' ENTERED AT 09:38:41 ON 06 APR 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the

the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 6 Apr 2006 VOL 144 ISS 15
FILE LAST UPDATED: 4 Apr 2006 (20060404/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 13

L4

8 L3

=> d 14 ibib abs tot

L4 ANSWER 1 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:1265191 HCAPLUS

DOCUMENT NUMBER: 144:22922

TITLE: Preparation of imidazolidinones for treatment of enteroviruses.

INVENTOR(S): Chern, Jyh-Haur; Shih, Shin-Ru; Chang, Chih-Shiang; Lee, Chung-Chi; Lee, Yen-Chun

PATENT ASSIGNEE(S): National Health Research Institutes, Taiwan

SOURCE: U.S. Pat. Appl. Publ., 22 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--------|-----------|-----------------|------------|
| US 2005267164 | A1 | 20051201 | US 2005-134936 | 20050523 |
| PRIORITY APPLN. INFO.: | | | US 2004-574266P | P 20040525 |
| OTHER SOURCE(S): | MARPAT | 144:22922 | | |

GI



AB Title compds. [I; R1 = H, halo, cyano, NO22, amino, alkyl, cycloalkyl, alkoxy, aryl, aralkyl, heterocycloalkyl, heteroaryl; R2 = H, alkyl,

cycloalkyl, aryl, aralkyl, heterocycloalkyl, heteroaryl; A1, A2 = aryl, aralkyl, heteroaryl; X, Y = CH₂, CHRa, CRaRb, NRc, O, S, SO, SO₂, aryl, cycloalkyl heterocyclyl, heteroaryl, alkenyl, alkynyl; Ra, Rb = halo, amino, alkyl, hydroxyalkyl, alkoxy, SH, alkylthio, aryl, aralkyl, heteroaryl; Rc = alkyl, aryl, aralkyl, cycloalkyl, heteroaryl, heterocycloalkyl; m, n, p = 0-5; x, y = 0, 1; Q = (CH₂)_mXx(CH₂)_nYy(CH₂)_pOA₂CH:NR₂; with a proviso], were prepared Thus, 1-(4-pyridyl)-2-imidazolidinone was stirred with NaH in DMF at 0°-room temperature followed by addition of 4-(5-bromopentyloxy)benzaldehyde O-Me oxime in DMF and stirring for 8 h to give 85% title compound (II). Several I showed IC₅₀ ≤38.1 nM against enterovirus EV71-2231 and EV71-4643 in vero cell monolayers.

L4 ANSWER 2 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:921409 HCAPLUS

DOCUMENT NUMBER: 143:386968

TITLE: Synthesis and antipicornavirus activity of (R)- and (S)-1-[5-(4'-chlorobiphenyl-4-yloxy)-3-methylpentyl]-3-pyridin-4-yl-imidazolidin-2-one

AUTHOR(S): Chern, Jyh-Haur; Chang, Chih-Shiang; Tai, Chia-Liang; ~~Lee, Yen-Chun~~; Lee, Chung-Chi; Kang, Iou-Jiun; Lee, Ching-Yin; Shih, Shin-Ru

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Zhunan Town, Taichung, Miaoli County, 350, Taiwan

SOURCE: Bioorganic & Medicinal Chemistry Letters (2005), 15(19), 4206-4211


CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB The new pyridyl imidazolidinone derivative, 1-[5-(4'-chlorobiphenyl-4-yloxy)-3-methylpentyl]-3-pyridin-4-ylimidazolidin-2-one [(±)-I], was synthesized and found to have an excellent antiviral activity against enterovirus 71 (EV71, IC₅₀ = 0.009 μM). Therefore, (S)-(+)-I and (R)-(-)-I were prepared starting from readily available monomethyl (R)-3-methylglutarate as a useful chiral building block and their antiviral activity was evaluated in a plaque reduction assay. Interestingly, (S)-(+)-I was 10-fold more active against EV71 (IC₅₀ = 0.003 μM) than (R)-(-)-I (IC₅₀ = 0.033 μM). Similar results were found against all five strains (1743, 2086, 2231, 4643, and BrCr) of EV71 tested. This demonstrated that the absolute

configuration of the chiral carbon atom at the 3-position of the alkyl linker considerably influenced the anti-EV71 activity of these pyridylimidazolidinones.

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:331777 HCAPLUS

DOCUMENT NUMBER: 143:43827

TITLE: Design, Synthesis, and Antipicornavirus Activity of 1-[5-(4-Arylphenoxy)alkyl]-3-pyridin-4-ylimidazolidin-2-one Derivatives

AUTHOR(S): Chang, Chih-Shiang; Lin, Ying-Ting; Shih, Shin-Ru; Lee, Chung-Chi; ~~Lee, Yen-Chun~~; Tai, Chia-Liang; Tseng, Sung-Nien; Chern, Jyh-Haur

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Zhunan, 350, Taiwan

SOURCE: Journal of Medicinal Chemistry (2005), 48(10), 3522-3535

CODEN: JMCMAR; ISSN: 0022-2625


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:43827

GI




AB A series of pyridylimidazolidinone derivs. was synthesized and tested in vitro against enterovirus 71 (EV71). On the basis of DBPR103 (I), introduction of a Me group at the 2- or 3-position of the linker between the imidazolidinone and the biphenyl resulted in markedly improved antiviral activity toward EV71 with IC50 values of 5.0 nM and 9.3 nM, resp. Increasing the branched chain to Pr resulted in a progressive decrease in activity, while inserting different heteroatoms entirely rendered the compound only weakly active. The introduction of a bulky group (cyclohexyl, Ph, or benzyl) led to loss of activity against EV71. The 4-chlorophenyl moiety was replaced with bioisosteric groups such as oxadiazole or tetrazole dramatically improving anti-EV71 activity and selectivity indexes. Some of these compds. exhibited a strong activity against lethal EV71, and no apparent cellular toxicity was observed. Three of the more potent imidazolidinone compds. were subjected to a large group of picornaviruses to determine their spectrum of antiviral activity.

REFERENCE COUNT: 96 THERE ARE 96 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2004:767274 HCAPLUS
DOCUMENT NUMBER: 141:410865
TITLE: Synthesis and antienteroviral activity of a series of novel, oxime ether-containing pyridyl imidazolidinones
AUTHOR(S): Chern, Jyh-Haur; Lee, Chung-Chi; Chang, Chih-Shiang; Lee, Yen-Chun; Tai, Chia-Liang; Lin, Ying-Ting; Shia, Kak-Shan; Lee, Ching-Yin; Shih, Shin-Ru
CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Taichung, 114, Taiwan
SOURCE: Bioorganic & Medicinal Chemistry Letters (2004), 14(20), 5051-5056
CODEN: BMCLE8; ISSN: 0960-894X
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 141:410865
GI



AB A series of oxime ether-containing pyridyl imidazolidinones, e.g., I, were synthesized and their antiviral activity was evaluated in a plaque reduction assay. This class of compds. was specific for human enteroviruses, in particular, enterovirus 71 (EV71). Some derivs. strongly inhibited enterovirus replication with activities higher or comparable to those of the reference compds. such as A1 and A2. Preliminary SAR studies revealed that the chain length of the alkyl linker and the alkyl substituent at the oxime ether group largely influenced the in vitro anti-EV71 activity of this class of potent antiviral agents. Among this series of compds. synthesized, the pyridyl imidazolidinone I, with an Et oxime ether group located at the para position of the phenoxy ring, was identified as the most potent enterovirus 71 inhibitor ($IC_{50} = 0.001 \mu M$) with no apparent cytotoxic effect toward RD (rhabdomyosarcoma) cell lines ($CC_{50} > 25 \mu M$). Furthermore, I has been shown broad-spectrum activity against most of the serotypes of enteroviruses tested in the nanomolar range.

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2004:743226 HCAPLUS
DOCUMENT NUMBER: 141:235774
TITLE: Mutation in enterovirus 71 capsid protein VP1 confers resistance to the inhibitory effects of pyridyl imidazolidinone
AUTHOR(S): Shih, Shin-Ru; Tsai, Mun-Chung; Tseng, Sung-Nien; Won, Kuo-Fang; Shia, Kak-Shan; Li, Wen-Tai; Chern,

Jyh-Haur; Chen, Guang-Wu; Lee, Chung-Chi; Lee, Yen-Chun; Peng, Kuan-Chang; Chao, Yu-Sheng
 CORPORATE SOURCE: School of Medical Technology, Chang Gung University, Taoyuan, Taiwan
 SOURCE: Antimicrobial Agents and Chemotherapy (2004), 48(9), 3523-3529
 CODEN: AMACCQ; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Enterovirus 71 is one of the most important pathogens in the family of Picornaviridae that can cause severe complications in the postpoliovirus era, such as encephalitis, pulmonary edema, and even death. Pyridyl imidazolidinone is a novel class of potent and selective human enterovirus 71 inhibitor. Pyridyl imidazolidinone was identified by using computer-assisted drug design. This virol. investigation demonstrates that BPR0Z-194, one of the pyridyl imidazolidinones, targets enterovirus 71 capsid protein VP1. Time course expts. revealed that BPR0Z-194 effectively inhibited virus replication in the early stages, implying that the compound can inhibit viral adsorption and/or viral RNA uncoating. BPR0Z-194 was used to select and characterize the drug-resistant viruses. Sequence anal. of the VP1 region showed that the resistant variants differed consistently by seven amino acids in VP1 region from their parental drug-sensitive strains. Site-directed mutagenesis of enterovirus 71 infectious cDNA revealed that a single amino acid alteration at the position 192 of VP1 can confer resistance to the inhibitory effects of BPR0Z-194.

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:490448 HCAPLUS

DOCUMENT NUMBER: 141:54337

TITLE: Preparation of imidazolidinones for treating enterovirus infection

INVENTOR(S): Chern, Jyh-Haur; Shih, Shin-Ru; Chen, Chiung-Tong; Chang, Chih-Shiang; Lee, Chung-Chi; Lee, Yen-Chun; ~~Taiwan~~ Chia-Liang

PATENT ASSIGNEE(S): Taiwan

SOURCE: U.S. Pat. Appl. Publ., 33 pp., Cont.-in-part of U.S. Ser. No. 191,941.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:


| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---------------|------|----------|-----------------|----------|
| US 2004116476 | A1 | 20040617 | US 2003-717786 | 20031119 |
| US 2003087936 | A1 | 20030508 | US 2002-191941 | 20020709 |
| US 6706739 | B2 | 20040316 | | |

PRIORITY APPLN. INFO.: US 2002-191941 A2 20020709
 US 2001-313878P P 20010821

OTHER SOURCE(S): MARPAT 141:54337
 GI



I



II

AB The title compds. [I; R1, R2 = H, halo, alkyl, aryl, etc.; A1, A2 = aryl, aralkyl, heteroaryl; X, Y = S, SO, substituted CH2, etc.; m, n, p = 0-5; x, y = 0-1 (at least one of x and y = 1); with provisos], useful in treating enterovirus infection, were prepared Thus, reacting 1-(4-pyridyl)-2-imidazolidinone with 4-(5-bromo-3-methylpentoxy)-4'-chlorobiphenyl in the presence of NaH in DMF afforded 71% II which showed antiviral activity against enterovirus, in particular, EV71, coxsackieviruses A9, and A24. The pharmaceutical composition comprising the compound I is claimed.

L4 ANSWER 7 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:355843 HCAPLUS

DOCUMENT NUMBER: 138:353994

TITLE: Preparation of substituted imidazolidinones as antiviral agents

INVENTOR(S): Shih, Kak-Shan; Shih, Shin-Ru; Chang, Chung-Ming; Chern, Jyh-Haur; Li, Wen-Tai; Chen, Shu-Jen; Hsu, Ming-Chu

PATENT ASSIGNEE(S): National Health Research Institute, Taiwan

SOURCE: U.S. Pat. Appl. Publ., 16 pp.

CODEN: USXXCO


DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|-------------------|----------|------------------|-------------|
| US 2003087936 | A1 | 20030508 | US 2002-191941 | 20020709 |
| US 6706739 | B2 | 20040616 | | |
| TW 589307 | B | 20040601 | TW 2002-91117489 | 20020802 |
| US 2004116476 | A1 | 20040617 | US 2003-717786 | 20031119 |
| PRIORITY APPLN. INFO.: | | | US 2001-313878P | P 20010821 |
| | | | US 2002-191941 | A2 20020709 |
| OTHER SOURCE(S): | MARPAT 138:353994 | | | |
| GI | | | | |



I



II

AB Title compds. I [R1, R3 = aryl, aralkyl, heteroaryl, etc.; R2 = H, alkyl, haloalkyl, aryl, etc.; T = NH, O; W = (CH2)1-40; m = 4-8; n, p = 0-1 provided at least one of n, p = 1] are prepared For instance, 1-(4-pyridyl)-2-imidazolidinone is reacted with 1-bromo-6-[4-(trifluoromethyl)phenoxy]hexane (DMF, NaH, 0°, 30 min) to give II. Selected compds. showed antiviral activity against enteroviruses, in particular, enterovirus 71 and coxsackieviruses A9 and A24.

L4 ANSWER 8 OF 8 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:177380 HCAPLUS

DOCUMENT NUMBER: 136:369658

TITLE: Design, Synthesis, and Structure-Activity Relationship of Pyridyl Imidazolidinones: A Novel Class of Potent and Selective Human Enterovirus 71 Inhibitors

AUTHOR(S): Shia, Kak-Shan; Li, Wen-Tai; Chang, Chung-Ming; Hsu, Ming-Chu; Chern, Jyh-Haur; Leong, Max K.; Tseng, Sung-Nien; Lee, Chung-Chi; Lee, Yen-Chun; Chen, Shu-Jen; Peng, Kuan-Chang; Tseng, Huan-Yi; Chang, Yi-Ling; Tai, Chia-Liang; Shih, Shin-Ru

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Taipei, 11529, Taiwan

SOURCE: Journal of Medicinal Chemistry (2002), 45(8), 1644-1655

CODEN: JMCMAR; ISSN: 0022-2623


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 136:369658

GI



AB When skeletons of Win compds. were used as templates, computer-assisted drug design led to the identification of a novel series of imidazolidinone derivs. with significant antiviral activity against enterovirus 71 (EV 71), the infection of which had resulted in about 80 fatalities during the 1998 epidemic outbreak in Taiwan. In addition to inhibiting all the genotypes (A, B, and C) of EV 71 in the submicromolar to low micromolar range, compds. I (R = Br, CF₃) were extensively evaluated against a variety of viruses, showing potent activity against coxsackievirus A9 (IC₅₀ = 0.47-0.55 μM) and coxsackievirus A24 (IC₅₀ = 0.47-0.55 μM) as well as moderate activity against enterovirus 68 (IC₅₀ = 2.13 μM) and echovirus 9 (IC₅₀ = 2.6 μM). Our SAR studies revealed that imidazolidinone analogs with an aryl substituent at the para position of the phenoxyl ring, such as II [R₁ = (un)substituted phenyl], in general exhibited the highest activity against EV 71. Among them, II (R₁ = Ph) and its hydrochloride salt, in terms of potency and selectivity index, appear to be the most promising candidates in this series for further development of anti-EV-71 agents. Preliminary results of the study on the mode of action by a time-course experiment suggest that test compds. I (R = Br, CF₃) can effectively inhibit virus replication in the early stages, referring to virus attachment or uncoating. This indicates that the surface protein may be the target for this type of compound

REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> FIL REGISTRY

| | | |
|--|------------------|---------------|
| COST IN U.S. DOLLARS | SINCE FILE ENTRY | TOTAL SESSION |
| FULL ESTIMATED COST | 39.63 | 207.22 |
| DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) | SINCE FILE ENTRY | TOTAL SESSION |
| CA SUBSCRIBER PRICE | -6.00 | -6.00 |

FILE 'REGISTRY' ENTERED AT 09:43:01 ON 06 APR 2006
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 4 APR 2006 HIGHEST RN 879269-14-4

04/06/2006 10717786.trn

DICTIONARY FILE UPDATES: 4 APR 2006 HIGHEST RN 879269-14-4

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

```
*****
*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*
*****
```


Structure search iteration limits have been increased. See HELP SLIMITS
for details.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10717786a.str



chain nodes :

6 7 8 16

ring nodes :

1 2 3 4 5 9 10 11 12 13 14

chain bonds :

1-6 2-7 5-16 7-8 8-10

ring bonds :

1-2 1-5 2-3 3-4 4-5 9-10 9-14 10-11 11-12 12-13 13-14

exact/norm bonds :

1-2 1-5 1-6 2-3 2-7 4-5 5-16 7-8 8-10

exact bonds :

3-4

normalized bonds :

9-10 9-14 10-11 11-12 12-13 13-14

isolated ring systems :

containing 1 : 9 :

04/06/2006 10717786.trn

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:CLASS 7:CLASS 8:CLASS 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 16:Atom

L5 STRUCTURE UPLOADED

=> d 15

L5 HAS NO ANSWERS

L5 STR



Structure attributes must be viewed using STN Express query preparation.

=> s 15

SAMPLE SEARCH INITIATED 09:43:43 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 7172 TO ITERATE

27.9% PROCESSED 2000 ITERATIONS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

1 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**

PROJECTED ITERATIONS: 138363 TO 148517
PROJECTED ANSWERS: 1 TO 184

L6 1 SEA SSS SAM L5

=> s 15 sss full

FULL SEARCH INITIATED 09:43:52 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 142527 TO ITERATE


100.0% PROCESSED 142527 ITERATIONS
SEARCH TIME: 00.00.05

157 ANSWERS

L7 157 SEA SSS FUL L5

=>

Uploading C:\Program Files\Stnexp\Queries\10717786b.str



chain nodes :
6 7 8 16
ring nodes :
1 2 3 4 5 9 10 11 12 13 14
chain bonds :
1-6 2-7 5-16 7-8 8-10
ring bonds :
1-2 1-5 2-3 3-4 4-5 9-10 9-14 10-11 11-12 12-13 13-14
exact/norm bonds :
1-2 1-5 1-6 2-3 2-7 4-5 5-16 7-8 8-10
exact bonds :
3-4
normalized bonds :
9-10 9-14 10-11 11-12 12-13 13-14
isolated ring systems :
containing 1 : 9 :

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:CLASS 7:CLASS 8:CLASS 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 16:Atom

L8 STRUCTURE UPLOADED

=> d 18

L8 HAS NO ANSWERS

L8 STR



Structure attributes must be viewed using STN Express query preparation.

04/06/2006 10717786.trn

=> s l8

SAMPLE SEARCH INITIATED 09:45:15 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 7172 TO ITERATE

27.9% PROCESSED 2000 ITERATIONS 4 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 138363 TO 148517
PROJECTED ANSWERS: 59 TO 513

L9 4 SEA SSS SAM L8

=> s l8 sss full

FULL SEARCH INITIATED 09:45:21 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 142527 TO ITERATE

100.0% PROCESSED 142527 ITERATIONS
SEARCH TIME: 00.00.05

286 ANSWERS

L10 286 SEA SSS FUL L8

=> FIL HCAPLUS

| | | |
|--|------------|---------|
| COST IN U.S. DOLLARS | SINCE FILE | TOTAL |
| | ENTRY | SESSION |
| FULL ESTIMATED COST | 334.76 | 541.98 |
| DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) | SINCE FILE | TOTAL |
| | ENTRY | SESSION |
| CA SUBSCRIBER PRICE | 0.00 | -6.00 |

FILE 'HCAPLUS' ENTERED AT 09:45:32 ON 06 APR 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 6 Apr 2006 VOL 144 ISS 15
FILE LAST UPDATED: 4 Apr 2006 (20060404/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

04/06/2006 10717786.trn

=> d his

(FILE 'HOME' ENTERED AT 09:37:04 ON 06 APR 2006)

FILE 'REGISTRY' ENTERED AT 09:37:13 ON 06 APR 2006

L1 STRUCTURE UPLOADED

L2 7 S L1

L3 132 S L1 SSS FULL

FILE 'HCAPLUS' ENTERED AT 09:38:41 ON 06 APR 2006

L4 8 S L3

FILE 'REGISTRY' ENTERED AT 09:43:01 ON 06 APR 2006

L5 STRUCTURE UPLOADED

L6 1 S L5

L7 157 S L5 SSS FULL

L8 STRUCTURE UPLOADED

L9 4 S L8

L10 286 S L8 SSS FULL

FILE 'HCAPLUS' ENTERED AT 09:45:32 ON 06 APR 2006

=> s 17

L11 9 L7

=> s 110

L12 21 L10

=> s 110 and py<=2002

21 L10

22795084 PY<=2002

L13 12 L10 AND PY<=2002

=> d 19 ibib abs tot

YOU HAVE REQUESTED DATA FROM FILE 'REGISTRY' - CONTINUE? (Y)/N:n

=> d 111 ibib abs tot

L11 ANSWER 1 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:1265191 HCAPLUS

DOCUMENT NUMBER: 144:22922

TITLE: Preparation of imidazolidinones for treatment of enteroviruses.

INVENTOR(S): Chern, Jyh-Haur; Shih, Shin-Ru; Chang, Chih-Shiang; Lee, Chung-Chi; Lee, Yen-Chun

PATENT ASSIGNEE(S): National Health Research Institutes, Taiwan

SOURCE: U.S. Pat. Appl. Publ., 22 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--------|------------|-----------------|------------|
| US 2005267164 | A1 | 2005-12-01 | US 2005-134936 | 20050523 |
| PRIORITY APPLN. INFO.: | | | US 2004-574266P | P 20040525 |
| OTHER SOURCE(S): | MARPAT | 144:22922 | | |

GI



AB Title compds. [I; R1 = H, halo, cyano, NO₂, amino, alkyl, cycloalkyl, alkoxy, aryl, aralkyl, heterocycloalkyl, heteroaryl; R2 = H, alkyl, cycloalkyl, aryl, aralkyl, heterocycloalkyl, heteroaryl; A1, A2 = aryl, aralkyl, heteroaryl; X, Y = CH₂, CHRa, CRaRb, NRc, O, S, SO, SO₂, aryl, cycloalkyl heterocyclyl, heteroaryl, alkenyl, alkynyl; Ra, Rb = halo, amino, alkyl, hydroxyalkyl, alkoxy, SH, alkylthio, aryl, aralkyl, heteroaryl; Rc = alkyl, aryl, aralkyl, cycloalkyl, heteroaryl, heterocycloalkyl; m, n, p = 0-5; x, y = 0, 1; Q = (CH₂)_mXx(CH₂)_nYy(CH₂)_pOA2CH:NOR2; with a proviso], were prepared. Thus, 1-(4-pyridyl)-2-imidazolidinone was stirred with NaH in DMF at 0°-room temperature followed by addition of 4-(5-bromopentyloxy)benzaldehyde O-Me oxime in DMF and stirring for 8 h to give 85% title compound (II). Several I showed IC₅₀ ≤ 38.1 nM against enterovirus EV71-2231 and EV71-4643 in *vero* cell monolayers.

L11 ANSWER 2 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:921409 HCAPLUS

DOCUMENT NUMBER: 143:386968

TITLE: Synthesis and antipicornavirus activity of (R)- and (S)-1-[5-(4'-chlorobiphenyl-4-yloxy)-3-methylpentyl]-3-pyridin-4-yl-imidazolidin-2-one

AUTHOR(S): ~~Chern~~, Jyh-Haur; Chang, Chih-Shiang; Tai, Chia-Liang; Lee, Yen-Chun; Lee, Chung-Chi; Kang, Iou-Jiun; Lee, Ching-Yin; Shih, Shin-Ru

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Zhunan Town, Taichung, Miaoli County, 350, Taiwan

SOURCE: Bioorganic & Medicinal Chemistry Letters (2005), 15(19), 4206-4211


CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB The new pyridyl imidazolidinone derivative, 1-[5-(4'-chlorobiphenyl-4-yloxy)-3-methylpentyl]-3-pyridin-4-ylimidazolidin-2-one [(±)-I], was synthesized and found to have an excellent antiviral activity against enterovirus 71 (EV71, $IC_{50} = 0.009 \mu M$). Therefore, (S)-(+)-I and (R)-(-)-I were prepared starting from readily available monomethyl (R)-3-methylglutarate as a useful chiral building block and their antiviral activity was evaluated in a plaque reduction assay. Interestingly, (S)-(+)-I was 10-fold more active against EV71 ($IC_{50} = 0.003 \mu M$) than (R)-(-)-I ($IC_{50} = 0.033 \mu M$). Similar results were found against all five strains (1743, 2086, 2231, 4643, and BrCr) of EV71 tested. This demonstrated that the absolute configuration of the chiral carbon atom at the 3-position of the alkyl linker considerably influenced the anti-EV71 activity of these pyridylimidazolidinones.

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:331777 HCAPLUS

DOCUMENT NUMBER: 143:43827

TITLE: Design, Synthesis, and Antipicornavirus Activity of 1-[5-(4-Arylphenoxy)alkyl]-3-pyridin-4-ylimidazolidin-2-one Derivatives

AUTHOR(S): Chang, ~~Chin-Shiang~~; Lin, Ying-Ting; Shih, Shin-Ru; Lee, Chung-Chi; Lee, Yen-Chun; Tai, Chia-Liang; Tseng, Sung-Nien; Chern, Jyh-Haur

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Zhunan, 350, Taiwan

SOURCE: Journal of Medicinal Chemistry (2005), 48(10), 3522-3535

CODEN: JMCMAR; ISSN: 0022-2623


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:43827

GI



AB A series of pyridylimidazolidinone derivs. was synthesized and tested in vitro against enterovirus 71 (EV71). On the basis of DBPR103 (I), introduction of a Me group at the 2- or 3-position of the linker between the imidazolidinone and the biphenyl resulted in markedly improved antiviral activity toward EV71 with IC50 values of 5.0 nM and 9.3 nM, resp. Increasing the branched chain to Pr resulted in a progressive decrease in activity, while inserting different heteroatoms entirely rendered the compound only weakly active. The introduction of a bulky group (cyclohexyl, Ph, or benzyl) led to loss of activity against EV71. The 4-chlorophenyl moiety was replaced with bioisosteric groups such as oxadiazole or tetrazole dramatically improving anti-EV71 activity and selectivity indexes. Some of these compds. exhibited a strong activity against lethal EV71, and no apparent cellular toxicity was observed. Three of the more potent imidazolidinone compds. were subjected to a large group of picornaviruses to determine their spectrum of antiviral activity.

REFERENCE COUNT: 96 THERE ARE 96 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 4 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:767274 HCAPLUS

DOCUMENT NUMBER: 141:410865

TITLE: Synthesis and antienteroviral activity of a series of novel, oxime ether-containing pyridyl imidazolidinones
 AUTHOR(S): Chern, Jyh-Haur; Lee, Chung-Chi; Chang, Chih-Shiang; Lee, Yen-Chun; Tai, Chia-Liang; Lin, Ying-Ting; Shia, Kak-Shan; Lee, Ching-Yin; Shih, Shin-Ru

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Taichung, 114, Taiwan

SOURCE: Bioorganic & Medicinal Chemistry Letters (2004), 14(20), 5051-5056

CODEN: BMCLE8; ISSN: 0960-894X


PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 141:410865

GI



AB A series of oxime ether-containing pyridyl imidazolidinones, e.g., I, were synthesized and their antiviral activity was evaluated in a plaque reduction assay. This class of compds. was specific for human enteroviruses, in particular, enterovirus 71 (EV71). Some derivs. strongly inhibited enterovirus replication with activities higher or comparable to those of the reference compds. such as A1 and A2. Preliminary SAR studies revealed that the chain length of the alkyl linker and the alkyl substituent at the oxime ether group largely influenced the in vitro anti-EV71 activity of this class of potent antiviral agents. Among this series of compds. synthesized, the pyridyl imidazolidinone I, with an Et oxime ether group located at the para position of the phenoxy ring, was identified as the most potent enterovirus 71 inhibitor (IC₅₀ = 0.001 μ M) with no apparent cytotoxic effect toward RD (rhabdomyosarcoma) cell lines (CC₅₀ > 25 μ M). Furthermore, I has been shown broad-spectrum activity against most of the serotypes of enteroviruses tested in the nanomolar range.

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 5 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:743226 HCAPLUS

DOCUMENT NUMBER: 141:235774

TITLE: Mutation in enterovirus 71 capsid protein VP1 confers resistance to the inhibitory effects of pyridyl imidazolidinone

AUTHOR(S): Shih, Shin-Ru; Tsai, Mun-Chung; Tseng, Sung-Nien; Won, Kuo-Fang; Shia, Kak-Shan; Li, Wen-Tai; Chern, Jyh-Haur; Chen, Guang-Wu; Lee, Chung-Chi; Lee, Yen-Chun; Peng, Kuan-Chang; Chao, Yu-Sheng

CORPORATE SOURCE: School of Medical Technology, Chang Gung University, Taoyuan, Taiwan

SOURCE: Antimicrobial Agents and Chemotherapy (2004), 48(9), 3523-3529

CODEN: AMACQ; ISSN: 0066-4804

PUBLISHER: American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Enterovirus 71 is one of the most important pathogens in the family of Picornaviridae that can cause severe complications in the postpoliovirus era, such as encephalitis, pulmonary edema, and even death. Pyridyl imidazolidinone is a novel class of potent and selective human enterovirus 71 inhibitor. Pyridyl imidazolidinone was identified by using computer-assisted drug design. This virol. investigation demonstrates that BPR0Z-194, one of the pyridyl imidazolidinones, targets enterovirus 71 capsid protein VP1. Time course expts. revealed that BPR0Z-194 effectively inhibited virus replication in the early stages, implying that the compound can inhibit viral adsorption and/or viral RNA uncoating. BPR0Z-194 was used to select and characterize the drug-resistant viruses. Sequence anal. of the VP1 region showed that the resistant variants

differed consistently by seven amino acids in VP1 region from their parental drug-sensitive strains. Site-directed mutagenesis of enterovirus 71 infectious cDNA revealed that a single amino acid alteration at the position 192 of VP1 can confer resistance to the inhibitory effects of BPR0Z-194.

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 6 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:490448 HCAPLUS

DOCUMENT NUMBER: 141:54337

TITLE: Preparation of imidazolidinones for treating enterovirus infection

INVENTOR(S): Chern, Jyh-Haur; Shih, Shin-Ru; Chen, Chiung-Tong; Chang, Chih-Shiang; Lee, Chung-Chi; Lee, Yen-Chun; Tai, Chia-Liang

PATENT ASSIGNEE(S): Taiwan

SOURCE: U.S. Pat. Appl. Publ., 33 pp., Cont.-in-part of U.S. Ser. No. 191,941.

CODEN: USXXCO

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2


PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------------------|---------------------|-----------------|-------------|
| US 2004116476 | A1 | 20040617 | US 2003-717786 | 20031119 |
| US 2003087936 | A1 | 20030508 | US 2002-191941 | 20020709 |
| US 6706739 | B2 | 20040316 | | |
| PRIORITY APPLN. INFO.: | | | US 2002-191941 | A2 20020709 |
| | | | US 2001-313878P | P 20010821 |
| OTHER SOURCE(S): | MARPAT 141:54337 | | | |

GI



I



II

AB The title compds. [I; R1, R2 = H, halo, alkyl, aryl, etc.; A1, A2 = aryl,

aralkyl, heteroaryl; X, Y = S, SO, substituted CH₂, etc.; m, n, p = 0-5; x, y = 0-1 (at least one of x and y = 1); with provisos], useful in treating enterovirus infection, were prepared Thus, reacting 1-(4-pyridyl)-2-imidazolidinone with 4-(5-bromo-3-methylpentyloxy)-4'-chlorobiphenyl in the presence of NaH in DMF afforded 71% II which showed antiviral activity against enterovirus, in particular, EV71, coxsackieviruses A9, and A24. The pharmaceutical composition comprising the compound I is claimed.

L11 ANSWER 7 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:355843 HCAPLUS

DOCUMENT NUMBER: 138:353994

TITLE: Preparation of substituted imidazolidinones as antiviral agents

INVENTOR(S): Shia, Kak-Shan; Shih, Shin-Ru; Chang, Chung-Ming; Chern, Jyh-Haur; Li, Wen-Tai; Chen, Shu-Jen; Hsu, Ming-Chu

PATENT ASSIGNEE(S): National Health Research Institute, Taiwan

SOURCE: U.S. Pat. Appl. Publ., 16 pp.

CODEN: USXXCO



DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|---------------------|------------------|-------------|
| US 2003087936 | A1 | 20030508 | US 2002-191941 | 20020709 |
| US 6706739 | B2 | 20040316 | | |
| TW 589307 | B | 20040601 | TW 2002-91117489 | 20020802 |
| US 2004116476 | A1 | 20040617 | US 2003-717786 | 20031119 |
| PRIORITY APPLN. INFO.: | | | US 2001-313878P | P 20010821 |
| | | | US 2002-191941 | A2 20020709 |
| OTHER SOURCE(S): | | MARPAT 138:353994 | | |
| GI | | | | |



AB Title compds. I [R1, R3 = aryl, aralkyl, heteroaryl, etc.; R2 = H, alkyl, haloalkyl, aryl, etc.; T = NH, O; W = (CH₂)₁₋₄₀; m = 4-8; n, p = 0-1 provided at least one of n, p = 1] are prepared For instance, 1-(4-pyridyl)-2-imidazolidinone is reacted with 1-bromo-6-[4-(trifluoromethyl)phenoxy]hexane (DMF, NaH, 0°, 30 min) to give II. Selected compds. showed antiviral activity against enteroviruses, in particular, enterovirus 71 and coxsackieviruses A9 and A24.

L11 ANSWER 8 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:177380 HCAPLUS

DOCUMENT NUMBER: 136:369658

TITLE: Design, Synthesis, and Structure-Activity Relationship of Pyridyl Imidazolidinones: A Novel Class of Potent and Selective Human Enterovirus 71 Inhibitors

AUTHOR(S): Shia, Kak-Shan; Li, Wen-Tai; Chang, Chung-Ming; Hsu, Ming-Chu; Chern, Jyh-Haur; Leong, Max K.; Tseng, Sung-Nien; Lee, Chung-Chi; Lee, Yen-Chun; Chen, Shu-Jen; Peng, Kuan-Chang; Tseng, Huan-Yi; Chang, Yi-Ling; Tai, Chia-Liang; Shih, Shin-Ru

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Taipei, 11529, Taiwan

SOURCE: Journal of Medicinal Chemistry (2002), 45(8), 1644-1655

CODEN: JMCMAR; ISSN: 0022-2623



PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 136:369658

GI



AB When skeletons of Win compds. were used as templates, computer-assisted drug design led to the identification of a novel series of imidazolidinone derivs. with significant antiviral activity against enterovirus 71 (EV 71), the infection of which had resulted in about 80 fatalities during the 1998 epidemic outbreak in Taiwan. In addition to inhibiting all the genotypes (A, B, and C) of EV 71 in the submicromolar to low micromolar range, compds. I (R = Br, CF₃) were extensively evaluated against a variety of viruses, showing potent activity against coxsackievirus A9 (IC₅₀ = 0.47-0.55 μM) and coxsackievirus A24 (IC₅₀ = 0.47-0.55 μM) as well as moderate activity against enterovirus 68 (IC₅₀ = 2.13 μM)

and echovirus 9 (IC50 = 2.6 μ M). Our SAR studies revealed that imidazolidinone analogs with an aryl substituent at the para position of the phenoxy ring, such as II [R1 = (un)substituted phenyl], in general exhibited the highest activity against EV 71. Among them, II (R1 = Ph) and its hydrochloride salt, in terms of potency and selectivity index, appear to be the most promising candidates in this series for further development of anti-EV-71 agents. Preliminary results of the study on the mode of action by a time-course experiment suggest that test compds. I (R = Br, CF3) can effectively inhibit virus replication in the early stages, referring to virus attachment or uncoating. This indicates that the surface protein may be the target for this type of compound

REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 9 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1995:887878 HCAPLUS

DOCUMENT NUMBER: 123:286023

TITLE: Preparation of 5-[4-(heterocyclylalkoxy)benzyl - or benzylidene]thiazolidine-2,4-dione derivatives as hypolipidemics and hypoglycemics

INVENTOR(S): Yano, Shingo; Ogawa, Kazuo; Fukushima, Masakazu

PATENT ASSIGNEE(S): Taiho Pharmaceutical Co Ltd, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 57 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|-----------------|------------|
| JP 07138258 | A2 | 19950530 | JP 1993-286509 | 19931116 |
| CA 2177553 | AA | 19971129 | CA 1996-2177553 | 19960528 |
| PRIORITY APPLN. INFO.: | | | JP 1993-286509 | A 19931116 |

OTHER SOURCE(S): MARPAT 123:286023

GI For diagram(s), see printed CA Issue.

AB The title compds. [I; R1, R2 = H, halo, lower (halo)alkyl or (halo)alkoxy; or R1 and R2 are bonded together to form C1-3 alkylenedioxy; X = N, CH; the single bond with a dotted line represents a single bond or a double bond; A = heterocyclyl selected from Q - Q5; R3, R4 = H, lower alkyl; n = 1-4], having little side effects and useful as antidiabetics having activity for lowering both sugar and lipids in blood, are prepared Thus, a solution of benzaldehyde derivative Q6-CHO (R1 = CF3) (preparation given) 9.5, 2,4-thiazolidinone 3.8, and AcONa 4.3 g in 50 mL toluene was refluxed for 15 h and the solvent was removed by distillation to give, after treatment with 80% aqueous AcOH and filtration of precipitated crystals, 76%

5-benzylidene-2,4-

thiazolidinone derivative (II; R = Q6, wherein R1 = CF3) which was hydrogenated over 5% Pd-C in 1,4-dioxane at 50° and H pressure 50 atm to give 80% 5-benzyl-2,4-thiazolidinone derivative (III; R = Q6, wherein R1 = CF3) (IV). IV and III (R = Q6, wherein R1 = CF3) at 2.5 mg/kg p.o. twice a day for 5 consecutive days lowered the blood sugar level by 41 and 53%, resp., in mice.

=> d 113 ibib abs tot


L13 ANSWER 1 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:888730 HCAPLUS

DOCUMENT NUMBER: 137:384747
 TITLE: Preparation of arylsulfonylpyranhydroxamates as matrix metalloprotease and/or aggrecanase inhibitors
 INVENTOR(S): Barta, Thomas E.; Becker, Daniel P.; Bedell, Louis J.; Boehm, Terri L.; Fobian, Yvette M.; Freskos, John N.; Hockerman, Susan L.; Kassab, Darren J.; Kolodziej, Steve A.; McDonald, Joseph J.; Norton, Monica B.; Rico, Joseph G.; Talley, John J.; Villamil, Clara I.; Wang, Tijuan Jane
 PATENT ASSIGNEE(S): Pharmacia Corporation, USA
 SOURCE: PCT Int. Appl., 627 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|--------------|
| WO 2002092588 | A2 | 20021121 | WO 2002-US15257 | 20020510 <-- |
| WO 2002092588 | A3 | 20030227 | | |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2446586 | AA | 20021121 | CA 2002-2446586 | 20020510 <-- |
| EP 1385836 | A2 | 20040204 | EP 2002-729204 | 20020510 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR | | | | |
| BR 2002009525 | A | 20040309 | BR 2002-9525 | 20020510 |
| JP 2004530691 | T2 | 20041007 | JP 2002-589473 | 20020510 |
| ZA 2003008525 | A | 20050217 | ZA 2003-8525 | 20030131 |
| BG 108285 | A | 20040930 | BG 2003-108285 | 20031023 |
| NO 2003004995 | A | 20031216 | NO 2003-4995 | 20031110 |
| US 2005101641 | A1 | 20050512 | US 2004-992483 | 20041117 |
| PRIORITY APPLN. INFO.: | | | | |
| | | | US 2001-290375P | P 20010511 |
| | | | US 2002-142737 | A3 20020510 |
| | | | WO 2002-US15257 | W 20020510 |
| | | | US 2003-657034 | A3 20030905 |

OTHER SOURCE(S): MARPAT 137:384747
 GI



AB Title compds. [I; A1 = H, (substituted) alkylcarbonyl, alkoxycarbonyl, carbocyclylcarbonyl, heterocyclylcarbonyl, aminoalkylthiocarbonyl, etc.;

A2A3C = (substituted) heterocyclyl; E1 = O, S, SO, SO₂, NR₁, CONR₁, CR₁R₂; E2 = (substituted) alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl; E3 = CO, O₂C, CNR₃, NR₄, NR₄SO₂, S, SO, etc.; E4 = bond, (substituted) alkyl, alkenyl; E5 = H, OH, (substituted) alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, carbocyclyl, heterocyclyl; R₁, R₂ = H, (substituted) alkyl; with provisos], were prepared Thus, tetrahydro-4-[[4-[[5-(4-methoxyphenyl)-5-oxopentyl]oxy]phenyl]sulfonyl]-2H-pyran-4-carboxylic acid 1,1-dimethylethyl ester (preparation given) in CH₂Cl₂ was treated with Me₃SiCN and ZnI₂ to give 81% cyanohydrin. The product in DMF was treated with 1-hydroxybenzotriazole, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, N-methylmorpholine, and tetrahydropyranhydroxylamine to give 70% THP-protected hydroxamate. The latter was stirred with aqueous HCl in dioxane/MeOH to give 59% 4-[[4-[[[(4Z)-5-cyano-5-(4-methoxyphenyl)-4-pentenyl]oxy]phenyl]sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide. This inhibited MMP-13 with IC₅₀ = 0.2. nM.

L13 ANSWER 2 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:177380 HCAPLUS

DOCUMENT NUMBER: 136:369658

TITLE: Design, Synthesis, and Structure-Activity Relationship of Pyridyl Imidazolidinones: A Novel Class of Potent and Selective Human Enterovirus 71 Inhibitors

AUTHOR(S): Shia, Kak-Shan; Li, Wen-Tai; Chang, Chung-Ming; Hsu, Ming-Chu; Chern, Jyh-Haur; Leong, Max K.; Tseng, Sung-Nien; Lee, Chung-Chi; Lee, Yen-Chun; Chen, Shu-Jen; Peng, Kuan-Chang; Tseng, Huan-Yi; Chang, Yi-Ling; Tai, Chia-Liang; Shih, Shin-Ru

CORPORATE SOURCE: Division of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Taipei, 11529, Taiwan

SOURCE: Journal of Medicinal Chemistry (2002), 45(8), 1644-1655

CODEN: JMCMAR; ISSN: 0022-2623



PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 136:369658

GI



AB When skeletons of Win compds. were used as templates, computer-assisted

drug design led to the identification of a novel series of imidazolidinone derivs. with significant antiviral activity against enterovirus 71 (EV 71), the infection of which had resulted in about 80 fatalities during the 1998 epidemic outbreak in Taiwan. In addition to inhibiting all the genotypes (A, B, and C) of EV 71 in the submicromolar to low micromolar range, compds. I (R = Br, CF₃) were extensively evaluated against a variety of viruses, showing potent activity against coxsackievirus A9 (IC₅₀ = 0.47-0.55 μ M) and coxsackievirus A24 (IC₅₀ = 0.47-0.55 μ M) as well as moderate activity against enterovirus 68 (IC₅₀ = 2.13 μ M) and echovirus 9 (IC₅₀ = 2.6 μ M). Our SAR studies revealed that imidazolidinone analogs with an aryl substituent at the para position of the phenoxy ring, such as II [R₁ = (un)substituted phenyl], in general exhibited the highest activity against EV 71. Among them, II (R₁ = Ph) and its hydrochloride salt, in terms of potency and selectivity index, appear to be the most promising candidates in this series for further development of anti-EV-71 agents. Preliminary results of the study on the mode of action by a time-course experiment suggest that test compds. I (R = Br, CF₃) can effectively inhibit virus replication in the early stages, referring to virus attachment or uncoating. This indicates that the surface protein may be the target for this type of compound

REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 3 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1998:122836 HCAPLUS

DOCUMENT NUMBER: 128:204834

TITLE: Synthesis of 5-membered ring-type compounds as potential cholecystokinin receptor ligands

AUTHOR(S): Pentassuglia, Giorgio; Araldi, Gian Luca; Donati, Daniele; Feriani, Aldo; Oliosi, Beatrice; Pasquarello, Alessandra; Ursini, Antonella

CORPORATE SOURCE: Glaxo Wellcome S.p.A., Medicines Research Centre, Verona, 37135, Italy


SOURCE: Farmaco (1997), 52(10), 573-581
CODEN: FRMCE8; ISSN: 0014-827X

PUBLISHER: Societa Chimica Italiana

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB Imidazolidine-2,4-diones I (R = Ph, 2-ClC₆H₄, 1-adamantylmethyl, R' = H, 2-Cl, 4-Cl, 3,4-Cl₂, R'' = Ph, 2-naphthyl, PhO, etc.) and 1,5-di-Ph tetramic acid derivs. II (R = H, Cl) and III were selected in order to evaluate some 5-membered heterocyclic ring compds. as potential templates for the synthesis of CCK receptor ligands. All the compds. were evaluated in vitro towards both CCK-B and CCK-A receptors.

REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 4 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1995:887878 HCAPLUS

DOCUMENT NUMBER: 123:286023

TITLE: Preparation of 5-[4-(heterocyclylalkoxy)benzyl - or
benzylidene]thiazolidine-2,4-dione derivatives as
hypolipidemics ~~and hypoglycemics~~

INVENTOR(S): Yano, Shingo; Ogawa, Kazuo; Fukushima, Masakazu

PATENT ASSIGNEE(S) : Taiho Pharmaceutical Co Ltd, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 57 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|-----------------|--------------|
| JP 07138258 | A2 | 19950530 | JP 1993-286509 | 19931116 <-- |
| CA 2177553 | AA | 19971129 | CA 1996-2177553 | 19960528 <-- |
| SECURITY APPLN. INFO.: | | | JP 1993-286509 | A 19931116 |

OTHER SOURCE(S): MARPAT 123:286023

GI For diagram(s), see printed CA Issue.

AB The title compds. [I; R₁, R₂ = H, halo, lower (halo)alkyl or (halo)alkoxy; or R₁ and R₂ are bonded together to form C1-3 alkylenedioxy; X = N, CH; the single bond with a dotted line represents a single bond or a double bond; A = heterocyclcyl selected from Q - Q5; R₃, R₄ = H, lower alkyl; n =

1-4], having little side effects and useful as antidiabetics having activity for lowering both sugar and lipids in blood, are prepared Thus, a solution of benzaldehyde derivative Q6-CHO (R1 = CF3) (preparation given) 9.5, 2,4-thiazolidinone 3.8, and AcONa 4.3 g in 50 mL toluene was refluxed for 15 h and the solvent was removed by distillation to give, after treatment with 80% aqueous AcOH and filtration of precipitated crystals, 76%

5-benzylidene-2,4-


thiazolidinone derivative (II; R = Q6, wherein R1 = CF3) which was hydrogenated over 5% Pd-C in 1,4-dioxane at 50° and H pressure 50 atm to give 80% 5-benzyl-2,4-thiazolidinone derivative (III; R = Q6, wherein R1 = CF3) (IV).. IV and III (R = Q6, wherein R1 = CF3O) at 2.5 mg/kg p.o. twice a day for 5 consecutive days lowered the blood sugar level by 41 and 53%, resp., in mice.

L13 ANSWER 5 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1995:408786 HCAPLUS
DOCUMENT NUMBER: 122:290463
TITLE: Preparation of N-phenyl-N'-(phenoxyethyl)ureas or imidazolidinones as hypolipemics
INVENTOR(S): Ogawa, Kazuo; Oono, Tomoyasu; Yamada, Haruo
PATENT ASSIGNEE(S): Taiho Pharmaceutical Co Ltd, Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--------|------------|-----------------|--------------|
| JP 06345714 | A2 | 19941220 | JP 1993-140826 | 19930611 <-- |
| PRIORITY APPLN. INFO.: | | | JP 1993-140826 | 19930611 |
| OTHER SOURCE(S): | MARPAT | 122:290463 | | |

GI



AB The title compds. I [R1-2 = H, lower (halo)alkyl, lower (halo)alkoxy, halo; R3 = H, lower alkyl; R4 = H and R5 = H, lower alkyl or R4R5 = CH2CH2] and their salts are claimed. I inhibit formation of fatty acids and cholesterol. 1-(4-Chlorophenyl)-2-imidazolidinone (preparation given) was treated with NaH in DMF at room temperature for 30 min and the reaction mixture was further treated with 4-ClCH2CH2OC6H4CO2Me (DMF/THF solution) under stirring at 0° for 30 min and at room temperature for 18 h to give I (R1 = R2 = H, R3 = Me, R4R5 = CH2CH2). IC50 values of this compound against formation of fatty acids and sterols by isolated rat liver cells were 8.55 and 9.62 μM, resp.

L13 ANSWER 6 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1995:242246 HCAPLUS
DOCUMENT NUMBER: 122:80736

TITLE: Regio- and stereoselective synthesis of gem-difluorocyclopropanes using 4-bromo-4,4-difluorocrotonate

AUTHOR(S): Taguchi, Takeo; Sasaki, Hirofumi; Shibuya, Akira; Morikawa, Tsutomu

CORPORATE SOURCE: Tokyo College Pharmacy, Tokyo, 192-03, Japan

SOURCE: Tetrahedron Letters (1994), 35(6), 913-16
CODEN: TELEAY; ISSN: 0040-4039


PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 122:80736

GI



AB Regio- and stereoselective formation of the functionalized gem-difluorocyclopropanes was achieved through the Michael addition of lithium enolates of esters or amides with 4-bromo-4,4-difluorocrotonate followed by the triethylborane-O₂ mediated intramol. substitution reaction. Thus Michael addition of BrCF₂CH:CHCO₂C₆H₂Me₃-2,4,6 with the lithium enolate of AcOBu-t followed by cyclization in the presence of Et₃N, O₂, and 1,3-dimethyl-2-imidazolidinone gave 71% gem-difluorocyclopropane I.

L13 ANSWER 7 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1992:234832 HCAPLUS

DOCUMENT NUMBER: 116:234832

TITLE: Effect of aggregation on the reactivity of dodecylammonium propionate in organic solvents. 1. Kinetic models for esterolysis reactions

AUTHOR(S): Valeur, Bernard; Monnier, Eric

CORPORATE SOURCE: Lab. Chim. Gen., Conservatoire Natl. Metiers, Paris, 75003, Fr.

SOURCE: Journal of Colloid and Interface Science (1992), 150(2), 473-85

CODEN: JCISA5; ISSN: 0021-9797

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Dodecylammonium propionate (DAP) is a surfactant that undergoes sequential self-association in organic solvents. Its reactivity as a nucleophilic agent depends on the aggregation state. The kinetics of esterolysis of three allophanic esters and p-nitrophenyl acetate were studied in benzene and 1,2-dichloroethane as a function of surfactant concentration. After demonstrating by FTIR expts. that the monomeric form of DAP consists of dodecylamine and propionic acid, various kinetics models were developed. Excellent agreement of the exptl. data was found with a model in which two rate

consts., k_1 (for the monomeric form) and k_a (for a surfactant mol. in an aggregate, whatever the size of the aggregate), are considered. Anal. of the data also provides the equilibrium constant K of self-association. The validity of the kinetic model is further supported by the fact that the value found for K in benzene is the same for the four esters and in agreement with the value reported in the literature (determined by vapor pressure osmometry). Furthermore, this agreement means that the self-association of a surfactant like DAP can be studied by chemical kinetics.

L13 ANSWER 8 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN


ACCESSION NUMBER: 1983:594223 HCAPLUS
 DOCUMENT NUMBER: 99:194223
 TITLE: Base-catalyzed hydrolysis and decarboxylation of
 allophanic esters in a water-acetonitrile medium:
 bifunctional catalysis by a base-solvent entity
 AUTHOR(S): Monnier, E.; Klæbe, A.; Perie, J. J.
 CORPORATE SOURCE: Univ. Paul Sabatier, Toulouse, 31062, Fr.
 SOURCE: Tetrahedron Letters (1983), 24(30), 3067-70
 CODEN: TELEAY; ISSN: 0040-4039
 DOCUMENT TYPE: Journal
 LANGUAGE: French

AB Base-catalyzed hydrolysis of allophanic esters in MeCN-H₂O mixts. shows a sharp increase in rate constant in the range of 0.1-0.3 M H₂O, with a maximum, which is interpreted as a balance between two desolvation terms, one concerning the nucleophile, the other the anionic transition state. At low H₂O content (2×10^{-2} M), a fast hydrolysis ($k_{exp} \approx 0.5$ s⁻¹ at 20°) of allophanic esters is observed due to catalysis by the enolate of acetamide. This species also catalyzes the decarboxylation step, likely behaving as a bifunctional catalyst.

L13 ANSWER 9 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN


ACCESSION NUMBER: 1981:30750 HCAPLUS
 DOCUMENT NUMBER: 94:30750
 TITLE: 1-(Alkoxy carbonyl)-3-phenyl-5-methylhydantoins
 PATENT ASSIGNEE(S): Kyowa Hakko Kogyo Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|----------|----------|-----------------|--------------|
| ----- | ---- | ----- | ----- | ----- |
| JP 55051068 | A2 | 19800414 | JP 1978-124331 | 19781009 <-- |
| JP 60019731 | B4 | 19850517 | | |
| PRIORITY APPLN. INFO.: | | | JP 1978-124331 | A 19781009 |
| OTHER SOURCE(S): | CASREACT | 94:30750 | | |
| GI | | | | |



AB Fourteen hydantoin I (R = Me, Et, Bu, n-hexyl, CH₂Ph, Ph; R₁ = H, 4-Cl, 3-F, 3,4-Cl₂, etc.; Z = O, S), having antiinflammatory, analgesic, and muscle-relaxant activities, were prepared by acylation with ClCO₂R. Thus, 0.1 mol DL-alanine and 0.1 mol NaOH in aqueous MeCN treated with 0.1 mol 3,4-dichlorophenyl isocyanate gave 90.9% N-(3,4-dichlorophenylcarbamoyl)-DL-alanine, which was refluxed with 5N HCl 5 h to give 98.7% 3-(3,4-dichlorophenyl)-5-methylhydantoin. This (15 mmol) was treated with 20 mmol ClCO₂Et and Et₃N in EtOAc to give 90.5% I (R = Et, R₁ = 3,4-Cl₂, Z = O). Its optical isomers were prepared with D- or L-alanine instead of DL-alanine.

L13 ANSWER 10 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1979:22920 HCAPLUS
 DOCUMENT NUMBER: 90:22920
 TITLE: Novel syntheses of hydantoin derivatives
 AUTHOR(S): Iwata, K.; Hara, S.
 CORPORATE SOURCE: Cent. Res. Inst., Teijin Ltd., Hino, Japan
 SOURCE: Journal of Heterocyclic Chemistry (1978),
 15(7), 1231-4
 CODEN: JHTCAD; ISSN: 0022-152X
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB N-substituted hydantoin derivs. I (R = H, Ph, Me, 3-, 4-HO₂CC₆H₄; R₁ = Ph, Bu, 4-ClC₆H₄, etc.) were prepared by the condensation of an α-amino acid derivative, a primary amine, and di-Ph carbonate.

L13 ANSWER 11 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1975:589326 HCAPLUS
 DOCUMENT NUMBER: 83:189326
 TITLE: Plant growth regulator containing
 imidazolidinetrione-carboxylic acid derivatives
 INVENTOR(S): Baerlicher, Toni; Ebert, Edith
 PATENT ASSIGNEE(S): Ciba-Geigy A.-G., Switz.
 SOURCE: Patentschrift (Switz.), 11 pp.
 CODEN: SWXXAS
 DOCUMENT TYPE: Patent

LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|-----------------|--------------|
| CH 563711 | A | 19750715 | CH 1972-3131 | 19720303 <-- |
| ES 401164 | A1 | 19751116 | ES 1971-401164 | 19710325 <-- |
| US 3847940 | A | 19741112 | US 1972-311276 | 19721201 <-- |
| PRIORITY APPLN. INFO.: | | | CH 1971-4514 | A 19710326 |
| | | | US 1971-164007 | A3 19710719 |
| | | | CH 1971-11683 | A 19710806 |
| | | | CH 1972-3131 | A 19720303 |

GI For diagram(s), see printed CA Issue.

AB Imidazolidinetrionecarboxylic acid derivs. I (R1 = alkyl, alkoxy carbonyl, Ph, etc., R2 = alkyl, allyl, benzyl, halobenzyl, haloalkyl, etc.; Y = O or S) and 2,4,5-trioxoimidazolidine derivs. II (R1 = alkyl, halophenyl, etc.) are plant growth regulators. Thus, 0.2% 1-methyl-2,4,5-trioxoimidazolidine-3-carboxylic acid isobutyl ester [40407-00-9] decreased the force needed for plucking the orange fruit.

L13 ANSWER 12 OF 12 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1973:4249 HCAPLUS

DOCUMENT NUMBER: 78:4249

TITLE: Fruit abscission regulating and plant senescence inhibiting 2,4,5-trioxoimidazolidine derivatives

INVENTOR(S): Baerlocher, Toni; Ebert, Edith

PATENT ASSIGNEE(S): Ciba-Geigy A.-G.

SOURCE: Ger. Offen., 49 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|-----------------|--------------|
| DE 2214448 | A | 19721012 | DE 1972-2214448 | 19720324 <-- |
| DE 2214448 | B2 | 19810521 | | |
| DE 2214448 | C3 | 19820325 | | |
| CH 552596 | A | 19740815 | CH 1971-4514 | 19710326 <-- |
| CH 570381 | A | 19751215 | CH 1971-11683 | 19710806 <-- |
| IL 39042 | A1 | 19760430 | IL 1972-39042 | 19720320 <-- |
| CA 987328 | A1 | 19760413 | CA 1972-137606 | 19720321 <-- |
| IT 953546 | A | 19730810 | IT 1972-22257 | 19720322 <-- |
| AU 7240290 | A1 | 19730927 | AU 1972-40290 | 19720322 <-- |
| BE 781182 | A1 | 19720925 | BE 1972-115515 | 19720324 <-- |
| NL 7204027 | A | 19720928 | NL 1972-4027 | 19720324 <-- |
| FR 2131604 | A5 | 19721110 | FR 1972-10422 | 19720324 <-- |
| ZA 7202013 | A | 19721227 | ZA 1972-2013 | 19720324 <-- |
| BR 7201757 | A0 | 19730503 | BR 1972-1757 | 19720324 <-- |
| DD 102051 | C | 19731212 | DD 1972-161793 | 19720324 <-- |
| US 3818031 | A | 19740618 | US 1972-237926 | 19720324 <-- |
| GB 1373556 | A | 19741113 | GB 1972-14083 | 19720324 <-- |
| JP 56005201 | B4 | 19810204 | JP 1972-30208 | 19720325 <-- |
| US 3847940 | A | 19741112 | US 1972-311276 | 19721201 <-- |
| PRIORITY APPLN. INFO.: | | | CH 1971-4514 | A 19710326 |
| | | | CH 1971-11683 | A 19710806 |
| | | | CH 1971-11653 | A 19710506 |

US 1971-164007

A3 19710719

GI For diagram(s), see printed CA Issue.

AB Hundred and fifty-five title compds. (I. e.g. Q = O or S; R1 = H, alkyl, CH₂CH₂SEt, CH₂CO₂Et, CH₂Ch:CH₂, CH₂C.tplbond.CH, cycloalkyl, Ph, tetrahydrofurfuryl; R2 = alkyl, CH₂CH:CH₂, CH₂C.tplbond.CH, cyclohexyl, Ph, furfuryl) were prepared from II by successive reaction with Et₃N and ClCo-QR₂. I (Q = O) were also prepared by reaction of R₁NHCONHCO₂R₂ (III) with ClCOCOCl and if R₁ = H addnl. with R₁Br (R₁ = H). I inhibited the senescence of cut roses and Sinapis alba. The plucking strength of oranges from trees sprayed with 0.2-0.4% I solns. 7 days before the crop was <1.2-p.9 kg, as compared with 8.5 kg for untreated trees. Thus, ClCOCOCl was added to EtNHCONH₂ at 10°, the mixture stirred 3 hr at room temperature and refluxed 3 hr to give II (R₁ = Et) (IV). IV reacted successively with Et₃N at 15 and with ClCO₂CH₂CHMe₂ at 5-10° to give I (R₁ = Et, R₂ = CH₂CHMe₂) (V). H₂NCO₂CH₂CHMe₂ in PhMe and then EtNCO were added to NaH in PhMe at -10 to 0 and -10 to +10°, resp., to give III (R₁ = Et, R₂ = CH₂CHMe₂) (VI). VI and ClCOCOCl in CHCl₃ were stirred 2 hr at room temperature and refluxed 1 hr to give V. Compds. containing I were reported.

=> log y

COST IN U.S. DOLLARS

| SINCE FILE | TOTAL |
|------------|---------|
| ENTRY | SESSION |
| 65.13 | 607.11 |

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

| SINCE FILE | TOTAL |
|------------|---------|
| ENTRY | SESSION |
| -15.75 | -21.75 |

CA SUBSCRIBER PRICE

STN INTERNATIONAL LOGOFF AT 09:47:32 ON 06 APR 2006